

ERRATUM

## **Erratum to: Transoral Resection of Human Papillomavirus (HPV)-Positive Squamous Cell Carcinoma of the Oropharynx: Outcomes with and Without Adjuvant Therapy**

**Ryan S. Jackson, MD<sup>1</sup>, Parul Sinha, MBBS, MS<sup>1</sup>, Joseph Zenga, MD<sup>1</sup>, Dorina Kallogjeri, MD, MPH<sup>1,2</sup>, Jasmina Suko, MSc<sup>1,2</sup>, Eliot Martin, PA-C<sup>3</sup>, Eric J. Moore, MD<sup>3</sup>, and Bruce H. Haughey, MBChB, MS<sup>4</sup>**

<sup>1</sup>Department of Otolaryngology-Head and Neck Surgery, Washington University School of Medicine, Campus Box 8115, St. Louis, MO; <sup>2</sup>Clinical Outcomes Research Office, Washington University School of Medicine, St. Louis, MO; <sup>3</sup>Department of Otolaryngology-Head and Neck Surgery, Mayo Clinic, Rochester, MN; <sup>4</sup>Head and Neck Surgery Center of Florida, Florida Hospital Celebration Health and Florida Hospital Cancer Institute, Celebration, FL

### **ERRATUM TO: ANN SURG ONCOL DOI 10.1245/S10434-017-6041-X**

In the last sentence of the third paragraph of the second page of the original article the word “not” was inadvertently omitted. Following is the corrected sentence:

We hypothesized that patients undergoing transoral surgery with neck dissection alone have a higher risk of

locoregional recurrence but that this does not have an impact on overall survival or disease-specific survival secondary to successful salvage.

---

The online version of the original article can be found under doi:[10.1245/s10434-017-6041-x](https://doi.org/10.1245/s10434-017-6041-x).

---

© Society of Surgical Oncology 2017

Published Online: 28 August 2017

R. S. Jackson, MD  
e-mail: [jackson.ryan@wust.edu](mailto:jackson.ryan@wust.edu)